

Antisymmetric tensor contribution to the muon $g - 2$

Terry Pilling¹ and Richard T. Hammond²
Department of Physics, North Dakota State University
Fargo, ND, 58105-5566

P.F. Kelly³
Department of Physics, Ave Maria College
Ypsilanti, MI, 48197

Abstract

We investigate the Kalb-Ramond antisymmetric tensor field as a solution to the muon $g - 2$ problem. In particular we calculate the lowest-order Kalb-Ramond contribution to the muon anomalous magnetic moment and find that we can fit the new experimental value for the anomaly by adjusting the coupling without affecting the electron anomalous magnetic moment results.

1 Introduction

Particle magnetic moments have provided a valuable test of QED and the Standard Model. In the magnetic moments of the electron and the muon we have the important situation that both the experimental measurements and the standard model predictions are extremely precise. Additional importance of the muon stems from the fact that it has a larger rest mass than the electron and hence is more sensitive to massive virtual particles and to new physics. See [1] for a review of the electron and the muon anomalous magnetic moments.

It was recently found that the complete standard model prediction for the muon magnetic moment differs from the experimentally determined value [2, 3] by 1.6 standard deviations⁴. For a nice summary of the experimental and theoretical analyses, as well as history and prospects, see [4]. The measured value is expressed as $a(\text{exp}) = 11659203 \pm 15 \times 10^{-10}$ (assuming CPT invariance) where $a = (g - 2)/2$. The standard model prediction is $a(\text{SM}) = 11659176.7 \pm 6.7 \times 10^{-10}$.

There are many possible extensions of the standard model which may contribute to the anomalous magnetic moment [5, 6]. For example, the minimal supersymmetric extension of the standard model (MSSM) [7, 8, 5, 9, 10, 11], higher dimensional Kaluza-Klein theories [12], as well as theories based on heretofore undiscovered interactions [6], are all of this type.

In the present paper we investigate a new interaction involving an antisymmetric tensor field which we present as a potential solution to the 1.6σ discrepancy.

2 The Antisymmetric Tensor Interaction

The Kalb-Ramond antisymmetric tensor field [13, 14] is encountered in string theory [15] and also arises in some theories of gravity with torsion [16]. The contribution of the antisymmetric tensor to the anomalous magnetic moment occurs at one-loop order due to the Kalb-Ramond-fermion vertex. The Feynman diagram for this process is shown in Fig. 1.

¹terry@offshell.phys.ndsu.nodak.edu

²rich.hammond@ndsu.nodak.edu

³pfkelly@avemaria.edu

⁴The discrepancy originally stated in [2] and discussed in [4, 26] turned out to be incorrect due to a sign error found in the part of the standard model calculation dealing with the pion pole contribution. We refer the interested reader to [3, 27, 28, 29, 30] and references therein for the details.




Figure 1: The Feynman diagram for the antisymmetric tensor contribution to the muon anomalous magnetic moment. We have written the photon polarization index σ , the antisymmetric tensor indices μ, ν, α, β , and the spinor indices s, s', r, r' . The initial and final muon momenta are p and p' , the momentum of the antisymmetric tensor field is $p - k$ and the momentum transferred to the photon is q .

Consider the following lagrangian [15, 17, 18]:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{12}H_{\mu\nu\lambda}H^{\mu\nu\lambda} + \bar{\psi} \left(i \not{\partial} - e \not{A} - m - \frac{g}{M} \sigma_{\mu\nu\lambda} H^{\mu\nu\lambda} \right) \psi. \quad (1)$$

The factor of M denotes a mass scale to make the coupling constant, g , dimensionless. We assume that the antisymmetric torsion, or Kalb-Ramond antisymmetric tensor field, $H^{\mu\nu\lambda}$, can be derived from a potential (i.e. $H = dB$ is exact)

$$H_{\mu\nu\lambda} = \partial_\mu B_{\nu\lambda} + \partial_\nu B_{\lambda\mu} + \partial_\lambda B_{\mu\nu}, \quad (2)$$

with $B_{\nu\lambda} = -B_{\lambda\nu}$, and

$$\sigma_{\alpha\beta\gamma} = i\epsilon_{\alpha\beta\gamma\mu}\gamma_5\gamma^\mu. \quad (3)$$

To find the propagator for the antisymmetric tensor field requires adding a gauge fixing term to the free lagrangian density

$$\begin{aligned} \mathcal{L}_0 + \mathcal{L}_{\text{gf}} &= -\frac{1}{12}H_{\mu\nu\lambda}H^{\mu\nu\lambda} - \frac{1}{4\alpha} (\partial_\mu B^{\mu\nu} \partial^\sigma B_{\sigma\nu} + \partial_\mu B^{\nu\mu} \partial^\sigma B_{\nu\sigma}) \\ &= \frac{1}{4}B^{\alpha\beta} \left[g_{\alpha\nu}g_{\beta\lambda} \square + g_{\beta\nu}\partial_\lambda\partial_\alpha + g_{\alpha\lambda}\partial_\nu\partial_\beta + \frac{1}{\alpha} (g_{\beta\lambda}\partial_\alpha\partial_\nu + g_{\alpha\nu}\partial_\beta\partial_\lambda) \right] B^{\nu\lambda}, \end{aligned} \quad (4)$$

where α is a gauge fixing parameter. We use the antisymmetry of $B_{\mu\nu}$ and substitute $i\partial_\mu \rightarrow k_\mu$ to arrive at the momentum space expression

$$\mathcal{L}_0 + \mathcal{L}_{\text{gf}} = -\frac{1}{4}B^{\alpha\beta} \left[g_{\alpha\nu}g_{\beta\lambda}k^2 + \left(\frac{1}{\alpha} - 1 \right) (g_{\beta\lambda}k_\alpha k_\nu + g_{\alpha\nu}k_\beta k_\lambda) \right] B^{\nu\lambda}. \quad (5)$$

Our gauge invariant propagator for the antisymmetric tensor field is the inverse of the quantity in brackets directly above and reads:

$$G^{\nu\alpha\lambda\beta} = \frac{1}{k^2} \left[g^{\nu\beta}g^{\lambda\alpha} - g^{\nu\alpha}g^{\lambda\beta} + \frac{(1-\alpha)}{k^2} (g^{\nu\alpha}k^\lambda k^\beta + g^{\lambda\beta}k^\nu k^\alpha - g^{\nu\beta}k^\lambda k^\alpha - g^{\lambda\alpha}k^\nu k^\beta) \right]. \quad (6)$$

With the gauge choice $\alpha = 1$ this reduces to

$$G^{\nu\alpha\lambda\beta} = \frac{1}{k^2} (g^{\nu\beta}g^{\lambda\alpha} - g^{\nu\alpha}g^{\lambda\beta}). \quad (7)$$

The vertex rule for the coupling of the antisymmetric tensor field to a spin- $\frac{1}{2}$ Dirac field is extracted from the interaction part of the lagrangian

$$\mathcal{L}_{\text{int}} = -\frac{g}{M} \bar{\psi} \sigma_{\mu\nu\lambda} H^{\mu\nu\lambda} \psi. \quad (8)$$

Substituting the definitions (2) into this gives

$$\begin{aligned} \mathcal{L}_{\text{int}} &= -\frac{g}{M} \bar{\psi} [(i\epsilon_{\mu\nu\lambda\sigma} \gamma_5 \gamma^\sigma) (\partial^\mu B^{\nu\lambda} + \partial^\nu B^{\lambda\mu} + \partial^\lambda B^{\mu\nu})] \psi \\ &\equiv -\frac{g}{M} \bar{\psi} \Lambda^{\alpha\beta} \psi B_{\alpha\beta}. \end{aligned} \quad (9)$$

Our vertex rule in momentum space is then seen to be

$$-\frac{g}{M} \Lambda^{\alpha\beta} = -\frac{g}{M} \epsilon_{\mu\nu\lambda\sigma} \gamma_5 \gamma^\sigma (g^{\nu\alpha} g^{\lambda\beta} k^\mu + g^{\lambda\alpha} g^{\mu\beta} k^\nu + g^{\mu\alpha} g^{\nu\beta} k^\lambda), \quad (10)$$

where the momentum k is that carried by the Kalb-Ramond field $B_{\mu\nu}$.

3 Fermion Anomalous Magnetic Moment

The corrections to the magnetic moment are due to exchanges of virtual particles which affect the measured value of the photon-fermion coupling. Here we specialize to the case of muons.

The QED prediction is (to 5 loops) [2, 3, 19]:

$$a(\text{QED}) = 116584705.7(2.9) \times 10^{-11}$$

which differs in the 5th significant figure from the experimental value. The electroweak corrections to the vertex amount to (to 2 loops) [2, 20, 21]:

$$a(\text{EW}) = 152(4) \times 10^{-11}$$

There is yet another standard model correction. It is when the first muon emits a photon which is energetic enough to produce quark anti-quark pairs which then annihilate into another photon absorbed by the final state muon. This hadronic correction to the moment is computed two separate ways – with results which are in agreement – (to 3 loops) [2, 22, 23]:

$$\begin{aligned} a(\text{hadron1}) &= 6739(67) \times 10^{-11} \\ \text{or} \quad a(\text{hadron2}) &= 6803(114) \times 10^{-11}. \end{aligned}$$

We must note that this is very difficult to calculate and that it contributes the greatest amount of uncertainty in the total standard model prediction.

So the complete standard model prediction is then:

$$\begin{aligned} a(\text{SM}) &= a(\text{QED}) + a(\text{EW}) + a(\text{hadron1}) = 116591597(67) \times 10^{-11} \\ \text{or (with } a(\text{hadron2})) &= 116591660(114) \times 10^{-11} \end{aligned} \quad (11)$$

We now see that this differs from experiment in the 6th digit, which is much better than the pure QED calculation.

The current standard model calculation differs from the latest experimental results and is outside of the experimental error bars. This motivates a search for sources beyond the standard model that may help to resolve the discrepancy.

4 Torsion contribution to the magnetic moment

We can use the Feynman rules derived in Section 2 to find the contribution of the antisymmetric tensor field to fermion anomalous magnetic moments. This may place a useful bound on the fermion-antisymmetric tensor coupling.

The vertex correction, shown in Fig. 1, is defined by $\Gamma^\mu = \gamma^\mu + \delta\Gamma^\mu$, where

$$\delta\Gamma^\sigma(p', p) = \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{g^2}{m^2} \Lambda_{s'r'}^{\alpha\beta}(p-k) [iG_{\alpha\beta\mu\nu}(p-k)] \Lambda_{rs}^{\mu\nu}(p-k) [iS_F(k)_{lr}] (-\gamma_{l'l}^\sigma) [iS_F(k')_{r'l'}] \right\}. \quad (12)$$

Inserting the expressions for the propagator (7) and vertex rule (10) and using the standard identities

$$\begin{aligned} \epsilon_{\mu\nu\alpha\beta} \epsilon^{\alpha\beta\lambda\sigma} &= -4\delta_{[\mu}^\lambda \delta_{\nu]}^\sigma \\ \gamma^\rho \not{k}' \gamma^\sigma \not{k} \gamma_\rho &= -2 \not{k} \gamma^\sigma \not{k}' + \epsilon \not{k}' \gamma^\sigma \not{k} \\ \gamma^\rho \gamma^\sigma \not{k} \gamma_\rho &= 4k^\sigma - \epsilon \gamma^\sigma \not{k} \\ \gamma^\rho \gamma^\sigma \gamma_\rho &= (\epsilon - 2) \gamma^\sigma \end{aligned} \quad (13)$$

our vertex correction becomes

$$\delta\Gamma^\sigma(p', p) = \frac{i36g^2}{M^2} \int \frac{d^4k}{(2\pi)^4} \left\{ \frac{A^\sigma}{[(k')^2 - m^2][k^2 - m^2]} - \frac{B^\sigma}{[k-p]^2[(k')^2 - m^2]} \right\}, \quad (14)$$

where

$$\begin{aligned} A^\sigma &= (8mk^\sigma + 4mq^\sigma - 2 \not{k} \gamma^\sigma \not{k} - 2 \not{k} \gamma^\sigma \not{q} - 2m^2 \gamma^\sigma) + \epsilon (\not{k} \gamma^\sigma \not{k} + \not{q} \gamma^\sigma \not{k} - 2mk^\sigma - m \not{q} \gamma^\sigma + m^2 \gamma^\sigma), \\ B^\sigma &= 2p \cdot k \gamma^\sigma + 2p \cdot q \gamma^\sigma - 2p^\sigma \not{k} - k^2 \gamma^\sigma - \not{k} \not{q} \gamma^\sigma + 2mp^\sigma - m^2 \gamma^\sigma. \end{aligned} \quad (15)$$

Power counting indicates a quadratic divergence in need of regularization. Feynman parametrization to combine denominator factors gives

$$\delta\Gamma^\sigma = \frac{i36g^2}{M^2} \int \frac{d^4k}{(2\pi)^4} \int_0^1 dx dy \delta(x+y-1) \left\{ \frac{N_1^\sigma + \epsilon N_2^\sigma}{[k^2 - \Delta_1]^2} - \frac{N_3^\sigma}{[k^2 - \Delta_2]^2} \right\}, \quad (16)$$

where

$$\begin{aligned} N_1^\sigma &= 4m(1-2x)q^\sigma - 2 \not{k} \gamma^\sigma \not{k} - 2xy(2m^2 + q^2) \gamma^\sigma - 2m^2 \gamma^\sigma \\ N_2^\sigma &= \not{k} \gamma^\sigma \not{k} - x^2 q^2 \gamma^\sigma - 2m^2 x^2 \gamma^\sigma - m(1-2x)q^\sigma - m^2 \gamma^\sigma + mQ^\sigma \\ N_3^\sigma &= (k^2 + m^2(1+x^2) + xq^2(1+x)) \gamma^\sigma \\ \Delta_1 &= m^2 - xyq^2 \\ \Delta_2 &= m^2 y^2 - 2xyq^2. \end{aligned} \quad (17)$$

We have used the fact that $Q \cdot q = (p+p') \cdot q = 0$. The terms containing q^σ will vanish after performing the x, y integrations. This is in accord with the Ward identity.

The only term that will contribute to the anomalous magnetic moment is the term in N_2^σ involving Q^σ . Extracting only this term we have

$$\delta\Gamma_{\text{anom.}}^\sigma = \frac{i36g^2}{M^2} \int \frac{d^4k}{(2\pi)^4} \int_0^1 dx dy \left\{ \frac{\delta(x+y-1) \epsilon m Q^\sigma}{[k^2 - \Delta_1]^2} \right\}. \quad (18)$$

Applying dimensional regularization, the k integration yields

$$\delta\Gamma_{\text{anom.}}^\sigma = \frac{-36g^2}{(4\pi)^2 M^2} \int_0^1 dx dy \delta(x+y-1) \left\{ \epsilon m Q^\sigma \left[\frac{2}{\epsilon} - \gamma + \ln \left(\frac{4\pi\mu^2}{\Delta} \right) \right] \right\}. \quad (19)$$

Keeping only the zeroth order term in ϵ we have

$$\delta\Gamma_{\text{anom.}}^\sigma = \frac{-72mg^2}{(4\pi)^2 M^2} Q^\sigma. \quad (20)$$

Using the Gordon identity the magnetic moment part is isolated

$$\delta\Gamma_{\text{anom.}}^\sigma = \frac{72img^2}{(4\pi)^2 M^2} \sigma^{\sigma\nu} q_\nu \equiv \frac{i\sigma^{\sigma\nu} q_\nu}{2m} F_2(q^2), \quad (21)$$

providing an explicit form of the structure function,

$$F_2(q^2) = \frac{144m^2}{M^2} \frac{g^2}{(4\pi)^2}. \quad (22)$$

Note that to this, the lowest, order $F_2(q^2)$ is independent of q^2 . The g -factor of the muon is defined by

$$\frac{g_\mu - 2}{2} = F_2(0) = \frac{9m_\mu^2 g^2}{M^2 \pi^2}. \quad (23)$$

This expression will allow us to set a bound on the value of the antisymmetric tensor coupling [24] by fitting to the experimental discrepancy for the muon g -factor.

The standard model prediction differs from experiment [3] by $\delta a = 25(16) \times 10^{-10}$. Using this difference as an upper bound to the antisymmetric tensor contribution we have

$$a_\mu(\text{antisym}) \leq 25 \times 10^{-10} \quad (24)$$

The muon mass has been measured [25] to be $m_\mu = 105.658$ MeV. Inserted into (23), this yields

$$\frac{g^2}{M^2} \leq 2.456 \times 10^{-7} \text{ GeV}^{-2} \simeq 2.5 \times 10^{-7} \text{ GeV}^{-2} \quad (25)$$

which provides an explicit numerical *upper* bound on the antisymmetric tensor coupling⁵. Notice that if g is of order 1, then the mass scale is set at $M \sim 2$ TeV which is 1) much smaller than the the Planck scale and 2) may be near the supersymmetry scale.

This simple coupling makes no distinction between the fermion families and hence it must have effect on the anomalous magnetic moments of the electron and the tau. The electron anomalous magnetic moment is given in [25] as

$$a_e = \frac{g_e - 2}{2} = 1159652187 \pm 4 \times 10^{-12}. \quad (26)$$

With our upper bound for the coupling, the Kalb-Ramond contribution to the electron's magnetic moment is

$$a_e(\text{antisym}) \lesssim 0.06 \times 10^{-12} \quad (27)$$

which is well within the experimental error. The experimental value for the tau anomaly is not very precisely known at present and the effect of the antisymmetric tensor interaction, while approximately 300 times larger than in the muon case, is expected to be unverifiable in the near future.

⁵References [16, 31] have used the expression $\mathcal{L}_{\text{int}} = -i\sqrt{\frac{\pi G}{12}} \kappa \bar{\psi} \sigma_{\mu\nu\lambda} H^{\mu\nu\lambda} \psi$ as their interaction lagrangian rather than the one that we have used in (1) above. In order to facilitate comparison with their results it should be noted that our bound on g^2/M^2 given in (25) translates into a bound of $\kappa \leq 1.19 \times 10^{16}$.

5 Conclusion

We have found that the interaction between a fermion and an antisymmetric tensor field such as the one arising in string theory and in Einstein-Cartan gravity can solve the problem with the muon anomalous magnetic moment without having a significant affect on the electron anomalous magnetic moment if the antisymmetric tensor coupling satisfies the bound (25).

There are many other possible contributions, beyond the standard model, to the muon anomalous magnetic moment. The most favoured candidates are supersymmetric partners to the standard model spectrum [9, 8, 11, 10]. If these other particles are indeed found then the bound on the Kalb-Ramond-fermion coupling would become stronger.

References

- [1] V. W. Hughes and T. Kinoshita, *Rev. Mod. Phys.* **71**, S133 (1999).
- [2] H. N. Brown *et. al.*, hep-ex/0102017; *Phys. Rev. Lett.* **86**, 2227 (2001).
- [3] D. W. Hertzog, hep-ex/0202024.
- [4] J. P. Miller, hep-ex/0111036.
- [5] A. Czarnecki and W. J. Marciano, *Phys. Rev.* **D64**, 013014 (2001).
- [6] M. B. Einhorn and J. Wudka, *Phys. Rev. Lett.* **87**, 071805 (2001).
- [7] J. Grifols and A. Méndez, *Phys. Rev.* **D26**, 1809 (1982).
- [8] J. L. Feng and K. T. Matchev, *Phys. Rev.* **D64**, 013014 (2001).
- [9] L. Everett, G. L. Kane, S. Rigolin, and L.-T. Wang, hep-ph/0102145.
- [10] J. Ellis, D. Nanopoulos, and K. A. Olive, hep-ph/0102331.
- [11] F. Ferrer and J. A. Grifols, *Phys. Rev.* **D56**, 7466 (1997).
- [12] X. Calmet and A. Neronov, hep-ph/0104278.
- [13] M. Kalb and P. Ramond, *Phys. Rev.* **D9**, 2273 (1974).
- [14] P. K. Townsend, *Physics Letters* **B88**, 97 (1979).
- [15] T. Pilling, Ph.D. Thesis, North Dakota State University, Fargo, ND. (2002).
- [16] R. T. Hammond, *Rep. Prog. Phys.* **65**, 599 (2002).
- [17] N. D. H. Dass and K. V. Shajesh, hep-th/0107006.
- [18] S. Miyamoto and T. Nakano, *Prog. of Theor. Phys.* **45**, 295 (1971).
- [19] T. Kinoshita, B. Nizic, and Y. Okamoto, *Phys. Rev.* **D41**, 593 (1990).
- [20] A. Czarnecki, B. Krause, and W. J. Marciano, *Phys. Rev.* **D52**, R2619 (1995).
- [21] A. Czarnecki, B. Krause, and W. J. Marciano, *Phys. Rev. Lett.* **76**, 3267 (1996).
- [22] T. Kinoshita, B. Nizic, and Y. Okamoto, *Phys. Rev.* **D31**, 2108 (1985).
- [23] J. Erler and M. Luo, *Phys. Rev. Lett.* **87**, 071804 (2001).
- [24] D. E. Neville, *Phys. Rev.* **D21**, 2075 (1980).

- [25] Particle Data Group, European Physical Journal **C15**, 81 (2000).
- [26] G. Onderwater, [hep-ex/0105036](#).
- [27] M. Hayakawa and T. Kinoshita, [hep-ph/0112102](#).
- [28] A. C. Ian Blokland and K. Melnikov, [hep-ph/0112117](#).
- [29] M. Knecht and A. Nyffeler, [hep-ph/0111058](#).
- [30] E. P. Johan Bijmans and J. Prades, [hep-ph/0112255](#).
- [31] R. T. Hammond, Phys. Rev. **D52**, 6918 (1995).